

Monopoles in Superloop Space

Mir Faizal¹ and Tsou Sheung Tsun²

¹Department of Physics and Astronomy,
University of Waterloo, Waterloo,
Ontario N2L 3G1, Canada

²Mathematical Institute, University of Oxford,
Andrew Wiles Building,
Radcliffe Observatory Quarter, Woodstock Road,
Oxford OX2 6GG, United Kingdom

Abstract

In this paper, we will analyse a four dimensional gauge theory with $\mathcal{N} = 1$ supersymmetry in superloop space formalism. We will thus obtain an expression for the connection in the infinite-dimensional superloop space. We will then use this connection to obtain an expression for the curvature of the infinite-dimensional superloop space. We will also show that this curvature is proportional to the Bianchi identity in spacetime. Thus, in absence of a monopole this curvature will vanish. However, it will not vanish if the superloop intersects the world-line of a monopole because the Bianchi will not hold at that point.

1 Introduction

Electric–magnetic duality in electromagnetism is an important physical concept which has helped us understand, both the role of this symmetry and topological concepts inherent in field theories. Since Dirac [1] we know that the existence of magnetic monopoles is equivalent to (electric) charge quantization which in turn is equivalent to the electromagnetic gauge group being compact (i. e. $U(1)$). In view of the crucial role of non-abelian gauge theories in modern-day particle physics, the question of a non-abelian version of duality and monopoles is much studied [2]–[3]. As it turns out, loop space is crucial to the study of these questions. In this paper, we attempt to put these questions in the context of superloops, as there are indications that some of the physical applications in ordinary loop space might be usefully applied in supersymmetric theories in various contexts to be given in more details below.

Monopoles in ordinary gauge theories have been studied using Polyakov loops (Dirac phase factor), which are closed loops in spacetime in which no trace is taken over the gauge group indices (unlike the Wilsons loops) [4]. Polyakov loop space has been generalized to $\mathcal{N} = 1$ superloop space in three dimensions [5]. It is interesting to further generalize this to $\mathcal{N} = 1$ superloop space in four dimensions, because then it can be used to study various important physical systems. The non-abelian generalization of the Hodge duality (which is exactly

the duality of abelian gauge theory in four dimensions) can only be constructed in loop space [6]-[7]. Moreover, it is proved that this generalized loop space duality does reduce to the Hodge duality in spacetime for an abelian gauge theory. Furthermore, the sources in the original theory, appear as monopoles in the dual theory, and monopoles in the original theory, become sources in the dual theory. So, this loop space duality can be used to construct a dual potential for the non-abelian gauge theories. It may be noted that this can not be done in three dimensions as the original Hodge duality does exist in three dimensions. Thus, we need a four dimensional superloop space to construct a dual potential for supersymmetric gauge theories. It may be noted that confinement problem in non-Abelian gauge theories has been discussed using 't Hooft's order-disorder parameters [8]. The explicit construction of these order-disorder parameters requires this dual potential constructed in the loop space [9]. So, four dimensional superloop space can be used to study these order-disorder parameters in supersymmetric gauge theories.

Furthermore, this dual potential has also been used for constructing a Dualized Standard Model [10]-[15]. Supersymmetry is also an important ingredient in construction of particle physics models beyond the Standard Model. It may be noted that the CKM matrix is not an identity matrix even in the supersymmetric Standard Models [16]-[17]. The results obtained in this paper can be used to explain this fact by constructing a supersymmetric Dualized Standard Model. In that supersymmetric Dualized Standard Model, there will exist a dual gauge symmetry apart from the usual gauge symmetry. Thus, in the supersymmetric Dualized Standard Model fermion generations will be expected to appear as dual colors. This symmetry is expected to be broken with interesting physical implications. Furthermore, with simple assumptions about the dual hypercharges of the dual Higgs fields, the fermion mass hierarchy and that the CKM matrix are expected to be just the identity at the tree level. However, the loop corrections are expected to give small but non-zero values to both to the lower generation fermion masses and the off-diagonal CKM matrix elements in the supersymmetric Dualized standard model. These results have already been obtained for the conventional Standard Model using the duality derived from the loop space formalism [18]-[19]. It will be interesting to obtain similar results for supersymmetric Standard Model using the present superloop formalism.

In this paper, we will construct four dimensional $\mathcal{N} = 1$ superloop space. From a supersymmetric point of view, this theory will have the same amount of supersymmetry as three dimensional theory with $\mathcal{N} = 2$ supersymmetry. So, the results of this paper can be used for analysing monopoles in three dimensional gauge theories, in $\mathcal{N} = 2$ superspace. It may be noted that in certain cases, monopoles in the ABJM theory can enhance its supersymmetry from $\mathcal{N} = 6$ supersymmetry to $\mathcal{N} = 8$ supersymmetry [20]-[21]. This is important as the theory of M2-branes is expected to have full $\mathcal{N} = 8$ supersymmetry. It is important to study the effect of non-abelian monopoles in the ABJM theory for this supersymmetry enhancement. Furthermore, it is expected that the study of monopoles in the ABJM theory can help us understand the physics behind multiple M5-branes [22]-[23]. The results of this paper can find direct applications for such analysis.

2 Superloop Variables

In this section we will construct Polyakov loops for four dimensional Yang-Mills theory with $\mathcal{N} = 1$ supersymmetry. Polyakov loops are holonomies of closed loops in spacetime, and they are thus defined via parametrized loops in spacetime. However, they are independent of the parametrization chosen. As (unlike Wilsons loops) no trace is taken over gauge group indices in a Polyakov loop, they are gauge group valued functionals in the infinite-dimensional loop space. Polyakov loops are sometimes called the Dirac phase factor in the physics literature. As Polyakov loops are gauge group valued functionals, only their logarithmic derivative can be defined. This logarithmic derivative is used to define a connection in this loop space. This connection is called the Polyakov variable and it measures the change in phase as one moves from one point in the loop space to a neighboring point [24, 25]. Furthermore, this connection can be used to construct a curvature tensor. This curvature tensor in the infinite-dimensional loop space is proportional to the Bianchi identity in spacetime, and so, it vanishes when ever the Bianchi identity is satisfied [26]. As the Bianchi identity is not satisfied in presence of a monopole, so it does not vanish if a monopole is present. A similar construction has been done for three dimensional dimensional Yang-Mills theory with $\mathcal{N} = 1$ supersymmetry [5], and in this section, we will generalize those results to a four dimensional Yang-Mills theory with $\mathcal{N} = 1$ supersymmetry.

A four dimensional Yang-Mills theory with $\mathcal{N} = 1$ supersymmetry can be parameterized by anticommuting coordinates θ^a and $\theta^{\dot{a}}$ along with the usual spacetime coordinates, $x_{a\dot{a}} = \sigma_{a\dot{a}}^\mu x_\mu$ [27]. Here $\sigma_{a\dot{a}}^\mu = (\sigma_{a\dot{a}}^0, -\sigma_{a\dot{a}}^i)$, with $\sigma_{a\dot{a}}^0$ being a two dimensional identity matrix and $\sigma_{a\dot{a}}^i$ being complex matrices with eigenvalues ± 1 called the Pauli matrices. The generators of $\mathcal{N} = 1$ supersymmetry satisfy $\{Q_a, Q_{\dot{a}}\} = -i\partial_{a\dot{a}}$, where $\partial_{a\dot{a}} = \sigma_{a\dot{a}}^\mu \partial_\mu$. It is useful to define the derivatives $D_a = \partial_a + i\theta^{\dot{a}}\partial_{a\dot{a}}$ and $D_{\dot{a}} = \partial_{\dot{a}} + i\theta^a\partial_{a\dot{a}}$, which commute with these generators of $\mathcal{N} = 1$ supersymmetry. They also satisfy, $\{D_a, D_{\dot{a}}\} = i\partial_{a\dot{a}}$.

Usually, a vector superfield, V , is used for constructing a four dimensional non-abelian gauge theory with $\mathcal{N} = 1$ supersymmetry. This vector superfield is a matrix valued superfield, $V = V^\alpha T_\alpha$, where $[T_\alpha, T_\beta] = if_{\alpha\beta}^\gamma T_\gamma$, and it transforms under gauge transformation as $\exp(V) \rightarrow \exp(i\bar{\Lambda}) \exp(V) \exp(-i\Lambda)$. It is possible to use a gauge called the Wess-Zumino gauge, where $[V]_| = [D_a V]_| = [D^a D_a V]_| = [D_{\dot{a}} V]_| = [D^{\dot{a}} D_{\dot{a}} V]_| = 0$, where $'|'$ means that the quantity is evaluated at $\theta = 0$. Now we can construct a covariant derivative, ∇_A from this vector superfield as

$$\begin{aligned}\nabla_A &= (-i\{\mathcal{D}_a, D_{\dot{a}}\}, \mathcal{D}_a, D_{\dot{a}}), \\ \exp(V)\nabla_A \exp(-V) &= (-i\{D_a, \mathcal{D}_{\dot{a}}\}, D_a, \mathcal{D}_{\dot{a}}),\end{aligned}\tag{1}$$

where $\mathcal{D}_a = \exp(-V)D_a \exp(V)$ and $\mathcal{D}_{\dot{a}} = \exp(V)D_{\dot{a}} \exp(-V)$. The transformation of this covariant derivative can now be written as $\exp(V)\nabla_A \exp(-V) \rightarrow \exp(i\bar{\Lambda}) \exp(V)\nabla_A \exp(-V) \exp(-i\bar{\Lambda})$, and $\nabla_A \rightarrow \exp(i\Lambda) \nabla_A \exp(-i\Lambda)$. Furthermore, from the explicit form of the covariant derivative, the following quantities also vanish, $F_{a\dot{a}} = F_{ab} = F_{\dot{a}\dot{b}} = 0$, and the remaining field strengths can be expressed in terms of a single spinor valued field strength, $2W_a = iD^{\dot{a}} D_{\dot{a}} \exp(-V)D_a \exp(V)$, where $W_{\dot{a}} = -\exp(-V)W_{\dot{a}}^\dagger \exp(V)$ and $\nabla^a W_a = -\nabla^{\dot{a}} W_{\dot{a}}$. Now we can view the derivatives $D_A = (\partial_{a\dot{a}}, D_a, D_{\dot{a}})$ as a super-

vector and thus define a super one-form as $\Gamma = \xi^{a\dot{a}}\Gamma_{a\dot{a}} + \xi^a\Gamma_a + \xi^{\dot{a}}\Gamma_{\dot{a}}$, where $\Gamma_{a\dot{a}} = \Gamma_{a\dot{a}}^\alpha T_\alpha$, $\Gamma_a = \Gamma_a^\alpha T_\alpha$, $\Gamma_{\dot{a}} = \Gamma_{\dot{a}}^\alpha T_\alpha$, and $[T_\alpha, T_\beta] = if_{\alpha\beta}^\gamma T_\gamma$. The covariant derivative can also be defined as $\nabla_A = D_A - i\Gamma_A$, such that $[\nabla_A, \nabla_B] = H_{AB}$, where $H_{AB} = T_{AB}^C \nabla_C - iF_{AB}$. Now the Bianchi identity can be written as $[\nabla_{[A}, H_{BC]}] = 0$. As we can express the field strength F_{AB} as $F_{AB} = D_{[A}\Gamma_{B]} - i[\Gamma_A, \Gamma_B] - T_{AB}^C \Gamma_C$, so, we can write $F_{a\dot{a}} = D_{[a}\Gamma_{\dot{a}]} + D_{[\dot{a}}\Gamma_{a]} - i[\Gamma_a, \Gamma_{\dot{a}}] - i\Gamma_{a\dot{a}}$. Now if we impose the constraint $F_{a\dot{a}} = 0$, we obtain, $i\Gamma_{a\dot{a}} = D_{[a}\Gamma_{\dot{a}]} + D_{[\dot{a}}\Gamma_{a]} - i[\Gamma_a, \Gamma_{\dot{a}}]$. Thus, by imposing the constraint, $F_{a\dot{a}} = 0$, we can express $\Gamma_{a\dot{a}}$ in terms of Γ_a and $\Gamma_{\dot{a}}$.

Now we will construct a loop space formalism for super-Yang-Mills theories in four dimensions. Thus, we first parameterizing the superloop space by the coordinates $\xi^A(s) = (\xi^{a\dot{a}}(s), \xi^a(s), \xi^{\dot{a}}(s))$,

$$C : \{\xi^A(s) : s = 0 \rightarrow 2\pi, \xi^A(0) = \xi^A(2\pi)\}, \quad (2)$$

where $\xi^A(0) = \xi^A(2\pi)$ is a fixed point in the superloop space. We now define a superloop space variable as,

$$\begin{aligned} \Phi[\xi] &= P_s \exp i \left(\int_0^{2\pi} \Gamma^{a\dot{a}}(\xi(s)) \frac{d\xi_{a\dot{a}}}{ds} + \Gamma^a(\xi(s)) \frac{d\xi_a}{ds} + \Gamma^{\dot{a}}(\xi(s)) \frac{d\xi_{\dot{a}}}{ds} \right) \\ &= P_s \exp i \int_0^{2\pi} \Gamma^A(\xi(s)) \frac{d\xi_A}{ds}. \end{aligned} \quad (3)$$

where P_s denotes ordering in s increasing from right to left. The derivative in s is taken from below. It may be noted that labeling the superloop variable by a fixed point is over complete since the superloop variable $\Phi[\xi]$, depends only on C and not on the manner in which it is parameterized. Thus, if we introduce another parameter $s' = f(s)$ instead of s , it will only change the variable in the integration and not its value. Thus, by using this new parameter there will be no change in the value of $\Phi[\xi]$. It may be noted that $\Phi[\xi]$ is a scalar superfield from the supersymmetric point of view. Thus, it is possible to project out various ordinary loop superfields from it.

Now as $\Phi[\xi]$ is a gauge group valued functional, we can only define its logarithmic derivative

$$F_A[\xi|s] = i\Phi^{-1}[\xi] \frac{\delta}{\delta \xi^A(s)} \Phi[\xi], \quad (4)$$

where $F_A[\xi|s] = (F_{a\dot{a}}[\xi|s], F_a[\xi|s], F_{\dot{a}}[\xi|s])$. This quantity is a supersymmetric generalization of the Polyakov variable. This acts like a connection in the superloop space. To see that we will first define a parallel transport from a point $\xi(s_1)$ to a point $\xi(s_2)$ as

$$\begin{aligned} \Phi[\xi : s_1, s_2] &= P_s \exp i \left(\int_{s_1}^{s_2} \Gamma^{a\dot{a}}(\xi(s)) \frac{d\xi_{a\dot{a}}}{ds} + \Gamma^a(\xi(s)) \frac{d\xi_a}{ds} + \Gamma^{\dot{a}}(\xi(s)) \frac{d\xi_{\dot{a}}}{ds} \right) \\ &= P_s \exp i \int_{s_1}^{s_2} \Gamma^A(\xi(s)) \frac{d\xi_A}{ds}. \end{aligned} \quad (5)$$

Now we will parallel transport from a fixed point along a fixed path to another point say s . After reaching s , we will take a detour then turn back along the same path till we reach the original point where we started from. Thus, the phase factor generated by going along the path from the original point to s

will be canceled by the phase factor generated by going from s to the original point. So, there will be no contribution from this path. However, there will be a contribution generated by the transport along the infinitesimal circuit at s . This contribution will be proportional to $H^{AB}(s)$,

$$F^A[\xi|s] = \Phi^{-1}[\xi : s, 0] H^{AB}(\xi(s)) \Phi^{-1}[\xi : s, 0] \frac{d\xi_B(s)}{ds}, \quad (6)$$

As $F^A[\xi|s]$ represents the change in phase of $\Phi[\xi]$ as one moves from one point in the superloop space to a neighboring point, we can regard it as a connection in the superloop space. It may be noted that even though $F^A[\xi|s]$ proportional to the field strength in spacetime, it can be viewed as a connection in the superloop space. Now we first define a covariant derivative in superloop space as

$$\nabla_A(s) = \frac{\delta}{\delta \xi^A(s)} + i F_A[\xi|s]. \quad (7)$$

Now the curvature $-i G_{AB}[\xi, s_1, s_2]$ of the loop space can be defined by taking a commutator of these two covariant derivatives, $[\nabla_A[\xi(s_1)], \nabla_B[\xi(s_2)]]$,

$$\begin{aligned} G_{AB}[\xi(s_1, s_2)] &= \frac{\delta}{\delta \xi^A(s_2)} F_B[\xi|s_1] - \frac{\delta}{\delta \xi^B(s_1)} F_A[\xi|s_2] \\ &\quad + i[F_A[\xi|s_1], F_B[\xi|s_2]]. \end{aligned} \quad (8)$$

3 Monopoles

In the previous section we constructed a curvature for the infinite-dimensional superloop space. In this section, we will use this curvature to analyse monopoles in the superloop space. In fact, it has been demonstrated that for a three dimensional Yang-Mills theory with $\mathcal{N} = 1$ supersymmetry, the superloop space curvature vanished in absence of a monopole [5]. Now we have prove a four dimensional generalization of that result and show that a four dimensional Yang-Mills theory with $\mathcal{N} = 1$ supersymmetry, the superloop space curvature will vanish in absence of a monopole. In order to calculate the curvature explicitly, we will calculate the functional derivative of $F^B[\xi|s_1]$ with respect to $\xi_A(s_2)$. This derivative can be calculated by taking two infinitesimal variation of $\Phi^{-1}[\xi_2]\Phi[\xi_3] - \Phi^{-1}[\xi]\Phi[\xi_1]$, where $\xi_3^A(s) = (\xi_3^{a\dot{a}}(s), \xi_3^a(s), \xi_3^{\dot{a}}(s)) = \xi_1^A(s) + \delta' \xi^A(s)$, $\xi_2^A(s) = (\xi_2^{a\dot{a}}(s), \xi_2^a(s), \xi_2^{\dot{a}}(s)) = \xi^A(s) + \delta' \xi^A(s)$, $\xi_1^A(s) = (\xi_1^{a\dot{a}}(s), \xi_1^a(s), \xi_1^{\dot{a}}(s)) = \xi^A(s) + \delta \xi^A(s)$, and $\delta \xi^A = \Delta^B \delta_B^A \delta(s - s')$, $\delta' \xi^A = \Delta'^B \delta_B^A \delta(s - s')$. Now we using parallel transport along these paths, we can write

$$\Phi[\xi_1] = \Phi[\xi] - i \int ds \Phi(\xi : 2\pi, s) \mathcal{H}(\xi(s)) \Phi(\xi : s, 0), \quad (9)$$

where

$$\begin{aligned} \mathcal{H}(\xi(s)) &= H^{a\dot{a}bb}(\xi(s)) \frac{d\xi_{bb}(s)}{ds} \delta \xi_{a\dot{a}}(s) + H^{ab\dot{b}}(\xi(s)) \frac{d\xi_{bb}(s)}{ds} \delta \xi_a(s) \\ &\quad + H^{\dot{a}bb}(\xi(s)) \frac{d\xi_{bb}(s)}{ds} \delta \xi_{\dot{a}}(s) + H^{a\dot{a}b}(\xi(s)) \frac{d\xi_b(s)}{ds} \delta \xi_{a\dot{a}}(s) \\ &\quad + H^{a\dot{a}\dot{b}}(\xi(s)) \frac{d\xi_{\dot{b}}(s)}{ds} \delta \xi_{a\dot{a}}(s) + H^{\dot{a}b}(\xi(s)) \frac{d\xi_b(s)}{ds} \delta \xi_{\dot{a}}(s) \end{aligned}$$

$$\begin{aligned}
& + H^{ab}(\xi(s)) \frac{d\xi_b(s)}{ds} \delta\xi_a(s) + H^{a\dot{b}}(\xi(s)) \frac{d\xi_{\dot{b}}(s)}{ds} \delta\xi_a(s) \\
& + H^{\dot{a}b}(\xi(s)) \frac{d\xi_b(s)}{ds} \delta\xi_{\dot{a}}(s).
\end{aligned} \tag{10}$$

Similarly, we can also write

$$\Phi[\xi_2] = \Phi[\xi] - i \int ds \Phi[\xi : 2\pi, s] \mathcal{H}(\xi'(s)) \Phi[\xi : s, 0], \tag{11}$$

where

$$\begin{aligned}
\mathcal{H}(\xi'(s)) = & H^{a\dot{a}b\dot{b}}(\xi(s)) \frac{d\xi_{b\dot{b}}(s)}{ds} \delta'\xi_{a\dot{a}}(s) + H^{ab\dot{b}}(\xi(s)) \frac{d\xi_{b\dot{b}}(s)}{ds} \delta'\xi_a(s) \\
& + H^{\dot{a}b\dot{b}}(\xi(s)) \frac{d\xi_{b\dot{b}}(s)}{ds} \delta'\xi_{\dot{a}}(s) + H^{a\dot{a}b}(\xi(s)) \frac{d\xi_b(s)}{ds} \delta'\xi_{a\dot{a}}(s) \\
& + H^{a\dot{a}\dot{b}}(\xi(s)) \frac{d\xi_{\dot{b}}(s)}{ds} \delta'\xi_{a\dot{a}}(s) + H^{\dot{a}b}(\xi(s)) \frac{d\xi_{\dot{b}}(s)}{ds} \delta'\xi_{\dot{a}}(s) \\
& + H^{ab}(\xi(s)) \frac{d\xi_b(s)}{ds} \delta'\xi_a(s) + H^{a\dot{b}}(\xi(s)) \frac{d\xi_{\dot{b}}(s)}{ds} \delta'\xi_a(s) \\
& + H^{\dot{a}b}(\xi(s)) \frac{d\xi_b(s)}{ds} \delta'\xi_{\dot{a}}(s).
\end{aligned} \tag{12}$$

Finally, we have

$$\Phi[\xi_3] = \Phi[\xi_1] - i \int ds \Phi[\xi_1 : 2\pi, s] \mathcal{H}(\xi_1(s)) \Phi[\xi_1 : s, 0], \tag{13}$$

where

$$\begin{aligned}
\mathcal{H}(\xi_1(s)) = & H^{a\dot{a}b\dot{b}}(\xi_1(s)) \frac{d\xi_{1b\dot{b}}(s)}{ds} \delta'\xi_{1a\dot{a}}(s) + H^{ab\dot{b}}(\xi_1(s)) \frac{d\xi_{1b\dot{b}}(s)}{ds} \delta'\xi_{1a}(s) \\
& + H^{\dot{a}b\dot{b}}(\xi_1(s)) \frac{d\xi_{1b\dot{b}}(s)}{ds} \delta'\xi_{1\dot{a}}(s) + H^{a\dot{a}b}(\xi_1(s)) \frac{d\xi_{1b}(s)}{ds} \delta'\xi_{1a\dot{a}}(s) \\
& + H^{a\dot{a}\dot{b}}(\xi_1(s)) \frac{d\xi_{1\dot{b}}(s)}{ds} \delta'\xi_{1a\dot{a}}(s) + H^{\dot{a}b}(\xi_1(s)) \frac{d\xi_{1\dot{b}}(s)}{ds} \delta'\xi_{1\dot{a}}(s) \\
& + H^{ab}(\xi_1(s)) \frac{d\xi_{1b}(s)}{ds} \delta'\xi_{1a}(s) + H^{a\dot{b}}(\xi_1(s)) \frac{d\xi_{1\dot{b}}(s)}{ds} \delta'\xi_{1a}(s) \\
& + H^{\dot{a}b}(\xi_1(s)) \frac{d\xi_{1b}(s)}{ds} \delta'\xi_{1\dot{a}}(s).
\end{aligned} \tag{14}$$

Here $\Phi[\xi_1 : s, 0]$ can be written as

$$\begin{aligned}
\Phi[\xi_1 : s, 0] = & \Phi[\xi : s, 0] - i \int_0^s ds' \Phi[\xi : s, s'] \mathcal{H}(\xi(s')) \Phi[\xi : s', 0] \\
& + i\Gamma^A(\xi(s)) \Phi[\xi : s, 0] \delta\xi_A(s),
\end{aligned} \tag{15}$$

where

$$\begin{aligned}
\mathcal{H}(\xi(s')) = & H^{a\dot{a}b\dot{b}}(\xi(s')) \frac{d\xi_{b\dot{b}}(s')}{ds'} \delta\xi_{a\dot{a}}(s') + H^{ab\dot{b}}(\xi(s')) \frac{d\xi_{b\dot{b}}(s')}{ds'} \delta\xi_a(s') \\
& + H^{\dot{a}b\dot{b}}(\xi(s')) \frac{d\xi_{b\dot{b}}(s')}{ds'} \delta\xi_{\dot{a}}(s') + H^{a\dot{a}b}(\xi(s')) \frac{d\xi_b(s')}{ds'} \delta\xi_{a\dot{a}}(s')
\end{aligned}$$

$$\begin{aligned}
& + H^{a\dot{a}\dot{b}}(\xi(s')) \frac{d\xi_{\dot{b}}(s')}{ds'} \delta\xi_{a\dot{a}}(s') + H^{\dot{a}\dot{b}}(\xi(s')) \frac{d\xi_{\dot{b}}(s')}{ds'} \delta\xi_{\dot{a}}(s') \\
& + H^{ab}(\xi(s')) \frac{d\xi_b(s')}{ds'} \delta\xi_a(s') + H^{a\dot{b}}(\xi(s')) \frac{d\xi_{\dot{b}}(s')}{ds'} \delta\xi_a(s') \\
& + H^{\dot{a}\dot{b}}(\xi(s')) \frac{d\xi_{\dot{b}}(s')}{ds'} \delta\xi_{\dot{a}}(s').
\end{aligned} \tag{16}$$

and a similar expression for $\Phi[\xi : 2\pi, s]$. Now collecting all the variations, we obtain the following expression,

$$\begin{aligned}
\frac{\delta}{\delta\xi_A(s_2)} F^B[\xi|s_1] &= \Phi^{-1}[\xi : s_1, 0] \nabla^B H^{AC}(\xi(s_2)) \\
&\times \frac{d\xi_C(s_1)}{ds_1} \Phi[\xi : s_1, 0] \delta(s_2 - s_1) \\
&+ \Phi^{-1}[\xi : s_2, 0] H_{AB}(\xi(s_2)) \Phi[\xi : s_2, 0] \\
&\times \frac{d}{ds_1} \delta(s_2 - s_1) \\
&+ i[F^A[\xi|s_2], F^B[\xi|s_1]] \theta(s_1 - s_2).
\end{aligned} \tag{17}$$

Similarly, we can write

$$\begin{aligned}
\frac{\delta}{\delta\xi_B(s_1)} F^A[\xi|s_2] &= \Phi^{-1}[\xi : s_2, 0] \nabla^A H^{BC}(\xi(s_1)) \\
&\times \frac{d\xi_C(s_2)}{ds_2} \Phi[\xi : s_2, 0] \delta(s_1 - s_2) \\
&+ \Phi^{-1}[\xi : s_1, 0] H_{BA}(\xi(s_1)) \Phi[\xi : s_1, 0] \\
&\times \frac{d}{ds_2} \delta(s_1 - s_2) \\
&+ i[F^B[\xi|s_1], F^A[\xi|s_2]] \theta(s_2 - s_1).
\end{aligned} \tag{18}$$

Now obtain an expression for the superloop space curvature,

$$\begin{aligned}
G_{AB}[\xi, s] &= \frac{\delta}{\delta\xi^A(s_2)} F_B[\xi|s_1] - \frac{\delta}{\delta\xi^B(s_1)} F_A[\xi|s_2] \\
&+ i[F_A[\xi|s_1], F_B[\xi|s_2]] \\
&= \Phi^{-1}[\xi : s, 0] [\nabla_{[A}, H_{BC)}] \Phi[\xi : s, 0] \frac{d\xi^C(s)}{ds}.
\end{aligned} \tag{19}$$

Thus, we observe that the superloop space curvature is proportional to the Bianchi identity in the spacetime. Now if the Bianchi identity are satisfied, $[\nabla_{[A}, H_{BC)}] = 0$, then the superloop space curvature will vanish, $G_{AB}[\xi, s] = 0$. The only non-vanishing contribution to the superloop space curvature can come from the existence of a monopole. This is because the Bianchi identity is not satisfied in presence of a monopole. So, if the world-line of a monopole intersects the superloop space, then the curvature of the superloop space will not vanish. In other words the non-vanishing of the superloop space curvature is an indicator for the existence of a monopole in spacetime.

4 Conclusion

In this paper we have analysed a four dimensional non-abelian gauge theory with $\mathcal{N} = 1$ supersymmetry in superloop space formalism. Thus, we were able to obtain an expression for a connection in this superloop space. An expression for the curvature of this superloop space was constructed using this connection. It was also demonstrated that this curvature is proportional to the Bianchi identity. Thus, it vanished if the Bianchi identity was satisfied. As the Bianchi identity is not satisfied in presence of a monopole, this curvature also did not vanish if the world-line of a monopole intersected the superloop space. In this way, the non-vanishing of the curvature acted as an indicator for the existence of a monopole in spacetime.

The results of this paper can be used for constructing a dual potential in supersymmetric gauge theories. This dual potential can be used for constructing a supersymmetric Dualized Standard Model. It can also be used for constructing 't Hooft's order-disorder parameters in supersymmetric gauge theories. The results obtained in this paper can also find important applications in M-theory. This is because they can be used for studying supersymmetry enhancement in the ABJM theory. They can also be used for understanding the physics of multiple M5-branes. It may be noted that D-branes in a graviphoton background [28]-[29], break the $\mathcal{N} = 1$ supersymmetry to $\mathcal{N} = 1/2$ supersymmetry [30]-[31]. It will be interesting to analyse monopoles in such a deformed field theory in four dimensions. In order to do so, we will have to construct a deformed version of the superloop space. This can possibly be done by replacing all the products of superfields with superstar products. It will be interesting to analyse the properties of monopoles in this deformed superspace.

References

- [1] P.A.M. Dirac, Proc. Roy. Soc. London A133, 60 (1931).
- [2] S. Coleman, Erice School, (1974)
- [3] T. T. Wu and C. N. Yang, Phys. Rev. D12, 3845 (1975).
- [4] A. M. Polyakov, Nucl. Phys. 164, 171 (1980).
- [5] M. Faizal, Europhys. Lett. 103, 21003 (2013).
- [6] H. M. Chan, J. Faridani and S. T. Tsou, Phys. Rev. D52, 6134 (1995)
- [7] H. M. Chan, J. Faridani and S. T. Tsou, Phys. Rev. D53 7293 (1996).
- [8] G. 't Hooft, Nucl. Phys. B138, 1 (1978).
- [9] H. M. Chan and S. T. Tsou, Phys. Rev. D56, 3646 (1997) .
- [10] H.M. Chan and S. T. Tsou, Phys. Rev. D57, 2507, (1998).
- [11] H. M. Chan and S. T. Tsou, Int. Jour. Mod. Phys. A14, 2173 (1999).
- [12] H. M. Chan and S. T. Tsou, Acta. Phys. Polon. B28, 3027 (1997).
- [13] H. M. Chan and S.T. Tsou, Acta. Phys. Polon. B33, 4041 (2002).

- [14] H. M. Chan, Int. J. Mod. Phys. A16, 163 (2001) .
- [15] H. M. Chan and S. T. Tsou, Acta. Phys. Polon. B28, 3041 (1997).
- [16] M. Kobayashi and T. Maskawa, Prog. Theor. Phys. 49, 652, (1973).
- [17] V. Barger, M. S. Berger and P. Ohmann, Phys. Rev. D 47, 2038 (1993).
- [18] J. Bordes, H. M. Chan, J. Faridani, J. Pfaudler and S. T. Tsou, Phys. Rev. D58, 013004 (1998).
- [19] J. Bordes , H. M. Chan, J. Faridani, J. Pfaudler and S.T. Tsou, Phys. Rev. D60, 013005 (1999).
- [20] O. K. Kwon, P. Oh and J. Sohn, JHEP. 0908, 093 (2009).
- [21] A. Gustavsson, JHEP. 1101, 037 (2011).
- [22] W. H. Huang, JHEP. 1105, 054 (2011).
- [23] S. Terashima and F. Yagi, JHEP. 1103, 036 (2011).
- [24] H. M. Chan, P. Scharbach and S. T. Tsou, Ann. Phys. 167, 454 (1986).
- [25] H. M. Chan and S. T. Tsou, Act. Phys. Pol. B17, 259 (1986).
- [26] H. M. Chan and S. T. Tsou, Some Elementary Gauge Theory Concepts, World Scientific, (1993).
- [27] S. J. Gates J, M.T. Grisaru, M. Rocek and W. Siegel, Front. Phys. 58, 1 (1983).
- [28] N. Seiberg, JHEP. 0306, 010 (2003).
- [29] N. Berkovits and N. Seiberg, JHEP. 0307, 010 (2003).
- [30] S. Ferrara, M. A. Lledo and O. Macia, JHEP. 0309, 068 (2003).
- [31] I. Bars, C. Deliduman, A. Pasqua and B. Zumino, Phys. Rev. D68, 106006 (2003).